



sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkylcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{19}$ or S);

5 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$, and

10 the dotted line indicates the presence of either a single or double bond, wherein the presence of a single bond, the valences are completed by hydrogens.


In a particular embodiment of the present invention, the compounds of the formula (II) are the following species:

 (II)								
A	B	D	R^1	R^2	R^3	R^4	R^5	R^6
O	O	O	Me	H	H	H	Me	Me
O	O	O	<i>i</i> -Pr	H	H	H	Me	Me
O	O	O	Ph	H	H	H	Me	Me
O	O	O	Me	Me	H	H	Me	Me
O	O	O	<i>i</i> -Pr	Me	H	H	Me	Me
O	O	O	Ph	Me	H	H	Me	Me
O	O	O	Me	H	Me	H	Me	Me
O	O	O	<i>i</i> -Pr	H	Me	H	Me	Me



(II)

A	B	D	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
O	O	O	Ph	H	Me	H	Me	Me
O	O	O	Me	H	H	Me	Me	Me
O	O	O	<i>i</i> -Pr	H	H	Me	Me	Me
O	O	O	Ph	H	H	Me	Me	Me
O	O	O	Me	H	CH ₂ Ph	H	Me	Me
O	O	O	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
O	O	O	Ph	H	CH ₂ Ph	H	Me	Me
O	CH ₂	O	Me	H	H	H	Me	Me
O	CH ₂	O	<i>i</i> -Pr	H	H	H	Me	Me
O	CH ₂	O	Ph	H	H	H	Me	Me
O	CH ₂	O	Me	Me	H	H	Me	Me
O	CH ₂	O	<i>i</i> -Pr	Me	H	H	Me	Me
O	CH ₂	O	Ph	Me	H	H	Me	Me
O	CH ₂	O	Me	H	Me	H	Me	Me
O	CH ₂	O	<i>i</i> -Pr	H	Me	H	Me	Me
O	CH ₂	O	Ph	H	Me	H	Me	Me
O	CH ₂	O	Me	H	H	Me	Me	Me
O	CH ₂	O	<i>i</i> -Pr	H	H	Me	Me	Me
O	CH ₂	O	Ph	H	H	Me	Me	Me



(II)

A	B	D	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
O	CH ₂	O	Me	H	CH ₂ Ph	H	Me	Me
O	CH ₂	O	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
O	CH ₂	CH ₂	Ph	H	CH ₂ Ph	H	Me	Me
O	CH ₂	CH ₂	Me	H	H	H	Me	Me
O	CH ₂	CH ₂	<i>i</i> -Pr	H	H	H	Me	Me
O	CH ₂	CH ₂	Ph	H	H	H	Me	Me
O	CH ₃	CH ₂	Me	Me	H	H	Me	Me
O	CH ₂	CH ₂	<i>i</i> -Pr	Me	H	H	Me	Me
O	CH ₂	CH ₂	Ph	Me	H	H	Me	Me
O	CH ₂	CH ₂	Me	H	Me	H	Me	Me
O	CH ₂	CH ₂	<i>i</i> -Pr	H	Me	H	Me	Me
O	CH ₂	CH ₂	Ph	H	Me	H	Me	Me
O	CH ₂	CH ₂	Me	H	H	Me	Me	Me
O	CH ₂	CH ₂	<i>i</i> -Pr	H	H	Me	Me	Me
O	CH ₂	CH ₂	Ph	H	H	Me	Me	Me
O	CH ₂	CH ₂	Me	H	CH ₂ Ph	H	Me	Me
O	CH ₂	CH ₂	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
O	CH ₂	CH ₂	Ph	H	CH ₂ Ph	H	Me	Me

In a sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O$, $B = O$, $E = O$ and $D = O$.

5 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O$, NR^{14} or S).

10 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}$ and R^{17} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O$, NR^{12} or S);

15 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

20 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O$, $B = NR^{20}$, $E = O$, $D = O$.

25 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O$, NR^{14} or S).

30 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}$ and R^{20} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

5 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}$, $\text{B} = \text{CR}^8\text{R}^9$, $\text{E} = \text{O}$, $\text{D} = \text{O}$.

10 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

15 R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

20 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

25 $\text{A} = \text{O}$, $\text{B} = \text{S}$, $\text{E} = \text{O}$, $\text{D} = \text{O}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$
and R^{22} independently are selected from the groups that include hydrogen, alkyl,
alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic,
sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,
phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,
carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or
 XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be
comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently
selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the
compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O, B = O, E = S, D = O.$

R^1 is selected independently from the groups that include hydrogen, alkyl,
cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,
a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$
or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$
and R^{22} independently are selected from the groups that include hydrogen, alkyl,
alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic,
sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,
phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,
carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or
 XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be
comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently
selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the
compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O, B = NR^{10}, E = S, D = O.$

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$ and R^{22} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


$A = O, B = CR^8R^9, E = S, D = O.$

20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$ and R^{22} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$ and R^{22} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$ and R^{22} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}$ and R^{22} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$
and R^{22} independently are selected from the groups that include hydrogen, alkyl,
alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic,
sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,
phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,
carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or
 XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be
comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently
selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the
compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl,
cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,
a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$
or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}$
and R^{22} independently are selected from the groups that include hydrogen, alkyl,
alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic,
sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,
phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,
carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or
 XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be
comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently
selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the
compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


$A = O, B = CR^8R^9, E = NR^{10}, D = O.$

20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



5 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

10 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

15 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}$, $\text{B} = \text{CR}^8\text{R}^9$, $\text{E} = \text{NR}^{10}$, $\text{D} = \text{O}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


$\text{A} = \text{CR}^8\text{R}^9$, $\text{B} = \text{O}$, $\text{E} = \text{O}$, $\text{D} = \text{O}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = CR^8R^9, B = S, E = O, D = O.$

20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^3 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.



In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{12}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.



In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,

phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

5 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




10 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

15 $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

20

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

25 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




30 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}$, $\text{B} = \text{O}$, $\text{E} = \text{S}$, $\text{D} = \text{CR}^8\text{R}^9$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


$\text{A} = \text{O}$, $\text{B} = \text{NR}^{10}$, $\text{E} = \text{S}$, $\text{D} = \text{CR}^8\text{R}^9$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

5 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}$, $\text{B} = \text{S}$, $\text{E} = \text{CR}^8\text{R}^9$, $\text{D} = \text{CR}^{20}\text{R}^{21}$.

10 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

15 R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

20 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

25 $\text{A} = \text{O}$, $\text{B} = \text{O}$, $\text{E} = \text{NR}^{16}$, $\text{D} = \text{CR}^8\text{R}^9$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O, B = NR^{10}, E = NR^{23}, D = CR^8R^9$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).


$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

$A = O, B = CR^7R^8$.

$E = NR^7, D = CR^7R^8$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

$\text{A} = \text{CR}^7\text{R}^8$, $\text{B} = \text{O}$.

$\text{E} = \text{O}$, $\text{D} = \text{CR}^7\text{R}^8$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{CR}^8\text{R}^9$, $\text{B} = \text{O}$, $\text{E} = \text{O}$, $\text{D} = \text{CR}^{20}\text{R}^{21}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{CR}^8\text{R}^9$, $\text{B} = \text{NR}^{10}$, $\text{E} = \text{O}$, $\text{D} = \text{CR}^{20}\text{R}^{21}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,

a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}, \text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}, \text{CR}^{15}=\text{CR}^{16}, \text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


$\text{A} = \text{CR}^8\text{R}^9, \text{B} = \text{CR}^{20}\text{R}^{21}, \text{E} = \text{O}, \text{D} = \text{CR}^{24}\text{R}^{25}$.

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}, \text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}, \text{CR}^{15}=\text{CR}^{16}, \text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

5 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{CR}^8\text{R}^9$, $\text{B} = \text{NR}^{10}$, $\text{E} = \text{S}$, $\text{D} = \text{CR}^{21}\text{R}^{22}$;

10 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

15 R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}$, NR^{12} or S);

20 R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:


25 $\text{A} = \text{CR}^8\text{R}^9$, $\text{B} = \text{CR}^{20}\text{R}^{21}$, $\text{E} = \text{S}$, $\text{D} = \text{CR}^{24}\text{R}^{25}$;

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}$, NR^{14} or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl,

carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

In another sub-embodiment, a structure of the formula (III) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:





R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S).

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);


10 R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$.

15 In a particular embodiment of the present invention, the compounds of the formula (III) are the following species:

 (III)								
B	D	E	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
O	O	O	Me	H	H	H	Me	Me
O	O	O	<i>i</i> -Pr	H	H	H	Me	Me
O	O	O	Ph	H	H	H	Me	Me
O	O	O	Me	Me	H	H	Me	Me
O	O	O	<i>i</i> -Pr	Me	H	H	Me	Me




B	D	E	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
O	O	O	Ph	Me	H	H	Me	Me
O	O	O	Me	H	Me	H	Me	Me
O	O	O	<i>i</i> -Pr	H	Me	H	Me	Me
O	O	O	Ph	H	Me	H	Me	Me
O	O	O	Me	H	H	Me	Me	Me
O	O	O	<i>i</i> -Pr	H	H	Me	Me	Me
O	O	O	Ph	H	H	Me	Me	Me
O	O	O	Me	H	CH ₂ Ph	H	Me	Me
O	O	O	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
O	O	O	Ph	H	CH ₂ Ph	H	Me	Me
CH ₂	O	O	Me	H	H	H	Me	Me
CH ₂	O	O	<i>i</i> -Pr	H	H	H	Me	Me
CH ₂	O	O	Ph	H	H	H	Me	Me
CH ₂	O	O	Me	Me	H	H	Me	Me
CH ₂	O	O	<i>i</i> -Pr	Me	H	H	Me	Me
CH ₂	O	O	Ph	Me	H	H	Me	Me
CH ₂	O	O	Me	H	Me	H	Me	Me
CH ₂	O	O	<i>i</i> -Pr	H	Me	H	Me	Me
CH ₂	O	O	Ph	H	Me	H	Me	Me



(III)

B	D	E	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
CH ₂	O	O	Me	H	H	Me	Me	Me
CH ₂	O	O	<i>i</i> -Pr	H	H	Me	Me	Me
CH ₂	O	O	Ph	H	H	Me	Me	Me
CH ₂	O	O	Me	H	CH ₂ Ph	H	Me	Me
CH ₂	O	O	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
CH ₂	O	O	Ph	H	CH ₂ Ph	H	Me	Me
CH ₂	CH ₂	O	Ph	H	CH ₂ Ph	H	Me	Me
CH ₂	CH ₂	O	Me	H	H	H	Me	Me
CH ₂	CH ₂	O	<i>i</i> -Pr	H	H	H	Me	Me
CH ₂	CH ₂	O	Ph	H	H	H	Me	Me
CH ₂	CH ₂	O	Me	Me	H	H	Me	Me
CH ₂	CH ₂	O	<i>i</i> -Pr	Me	H	H	Me	Me
CH ₂	CH ₂	O	Ph	Me	H	H	Me	Me
CH ₂	CH ₂	O	Me	H	Me	H	Me	Me
CH ₂	CH ₂	O	<i>i</i> -Pr	H	Me	H	Me	Me
CH ₂	CH ₂	O	Ph	H	Me	H	Me	Me
CH ₂	CH ₂	O	Me	H	H	Me	Me	Me
CH ₂	CH ₂	O	<i>i</i> -Pr	H	H	Me	Me	Me
CH ₂	CH ₂	O	Ph	H	H	Me	Me	Me



(III)

B	D	E	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶
CH ₂	CH ₂	O	Me	H	CH ₂ Ph	H	Me	Me
CH ₂	CH ₂	O	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
CH ₂	CH ₂	O	Ph	H	CH ₂ Ph	H	Me	Me
CH ₂	O	CH ₂	Me	H	H	H	Me	Me
CH ₂	O	CH ₂	<i>i</i> -Pr	H	H	H	Me	Me
CH ₂	O	CH ₂	Ph	H	H	H	Me	Me
CH ₂	O	CH ₂	Me	Me	H	H	Me	Me
CH ₂	O	CH ₂	<i>i</i> -Pr	Me	H	H	Me	Me
CH ₂	O	CH ₂	Ph	Me	H	H	Me	Me
CH ₂	O	CH ₂	Me	H	Me	H	Me	Me
CH ₂	O	CH ₂	<i>i</i> -Pr	H	Me	H	Me	Me
CH ₂	O	CH ₂	Ph	H	Me	H	Me	Me
CH ₂	O	CH ₂	Me	H	H	Me	Me	Me
CH ₂	O	CH ₂	<i>i</i> -Pr	H	H	Me	Me	Me
CH ₂	O	CH ₂	Ph	H	H	Me	Me	Me
CH ₂	O	CH ₂	Me	H	CH ₂ Ph	H	Me	Me
CH ₂	O	CH ₂	<i>i</i> -Pr	H	CH ₂ Ph	H	Me	Me
CH ₂	O	CH ₂	Ph	H	CH ₂ Ph	H	Me	Me

In a sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

A is O, S or NR⁷;

B and E are independently selected from CR⁸R⁹, O, S or NR¹⁰;

R¹ is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR⁷ (X = O, NR⁸ or S);

R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶ and R¹⁷ independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR¹¹ (X = O, NR¹² or S);

R¹ and R², R² and R³, R³ and R⁴, R⁴ and R⁵ and R⁵ and R⁶ can also each be comprised of one or two CR¹³R¹⁴ groups, connected by a tether, independently selected from CR¹⁵R¹⁶, CR¹⁵R¹⁶CR¹⁷R¹⁸, CR¹⁵=CR¹⁶, CR¹⁵R¹⁶O or CR¹⁵R¹⁶NR¹⁷;

and

the dotted line indicates the presence of either a single or double bond, wherein the presence of a single bond, the valences are completed by hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

A = O, B = O, E = O;


R¹ is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR¹³ (X = O, NR¹⁴ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

5 $A = O, B = CR^3R^9, E = O;$

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

10 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

20 the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O, B = S, E = O;$

25 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

30 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl,

alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}, \text{B} = \text{O}, \text{E} = \text{S};$


R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide,

phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}, \text{B} = \text{S}, \text{E} = \text{S};$

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}$, $\text{CR}^{15}=\text{CR}^{16}$, $\text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}; \text{B} = \text{O}, \text{E} = \text{CR}^8\text{R}^9;$

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

5 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}, CR^{15}R^{16}CR^{17}R^{18}, CR^{15}=CR^{16}, CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

15 the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O, B = NR^{10}, E = CR^7R^8$;

20 R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);


25 $R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

30

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,

a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{13}\text{R}^{14}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}, \text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}, \text{CR}^{15}=\text{CR}^{16}, \text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}; \text{B} = \text{O}, \text{E} = \text{NR}^{10}$;


R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:




R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O, NR^{14}$ or S);

$R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19}, R^{20}, R^{21}, R^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O, NR^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:



R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,

a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2, R^2 and R^3, R^3 and R^4, R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $\text{CR}^{15}\text{R}^{16}$ groups, connected by a tether, independently selected from $\text{CR}^{15}\text{R}^{16}, \text{CR}^{15}\text{R}^{16}\text{CR}^{17}\text{R}^{18}, \text{CR}^{15}=\text{CR}^{16}, \text{CR}^{15}\text{R}^{16}\text{O}$ or $\text{CR}^{15}\text{R}^{16}\text{NR}^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$\text{A} = \text{O}, \text{B} = \text{S}, \text{E} = \text{NR}^{10}$;

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($\text{X} = \text{O}, \text{NR}^{14}$ or S);

$\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{R}^{12}, \text{R}^{13}, \text{R}^{14}, \text{R}^{15}, \text{R}^{16}, \text{R}^{17}, \text{R}^{18}, \text{R}^{19}, \text{R}^{20}, \text{R}^{21}, \text{R}^{22}$ and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($\text{X} = \text{O}, \text{NR}^{12}$ or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = O$; $B = O$, $E = O$;

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{13} ($X = O$, NR^{14} or S);

R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} and R^{23} independently are selected from the groups that include hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, alkaryl, arylalkyl, heterocyclic, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, carboxylic acid, amide, nitro, cyano, azide, phosphonyl, phosphinyl, phosphoryl, phosphine, carbamate, ester, alkcarbonyl, carbonyl, halide, a residue of a natural or synthetic amino acid, or carbohydrate or XR^{11} ($X = O$, NR^{12} or S);

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^4 and R^5 and R^5 and R^6 can also each be comprised of one or two $CR^{13}R^{14}$ groups, connected by a tether, independently selected from $CR^{15}R^{16}$, $CR^{15}R^{16}CR^{17}R^{18}$, $CR^{15}=CR^{16}$, $CR^{15}R^{16}O$ or $CR^{15}R^{16}NR^{17}$;

the dotted line indicates the presence of either a single or double bond, wherein in the presence of a single bond, the valences are completed with hydrogens.

In another sub-embodiment, a structure of the formula (IV) is given wherein the compound or its pharmaceutically acceptable salts or prodrug is defined as follows:

$A = CR^8R^9$; $B = O$, $E = O$;

R^1 is selected independently from the groups that include hydrogen, alkyl, cycloalkyl, aryl, alkaryl, arylalkyl, heterocyclic, ester, alkcarbonyl, carbonyl, halide,